



Review

The Role of the Estrogen-Related Receptor Alpha (ERR α) in Hypoxia and Its Implications for Cancer Metabolism

Leslie Azarel Chaltel-Lima ^{1,2}, Fabiola Domínguez ³, Lenin Domínguez-Ramírez ^{3,*}
and Paulina Cortes-Hernandez ^{3,*}

¹ Segal Cancer Center, Lady Davis Institute for Medical Research, Jewish General Hospital, Montreal, QC, Canada; leslie.chaltellima@mail.mcgill.ca

² Division of Experimental Medicine, McGill University, Montreal, QC, Canada

³ Biología Celular, Centro de Investigación Biomédica de Oriente (CIBIOR), Instituto Mexicano del Seguro Social (IMSS), Atlixco, Pue, Mexico; irmafabiola@yahoo.com

* Correspondence: lenin.dominguez.ramierz@gmail.com (L.D.-R.); paulina.cortes.hernandez@gmail.com (P.C.-H.)

Abstract: Under low oxygen conditions (hypoxia), cells activate survival mechanisms including metabolic changes and angiogenesis, which are regulated by HIF-1. The estrogen-related receptor alpha (ERR α) is a transcription factor with important roles in the regulation of cellular metabolism that is overexpressed in hypoxia, suggesting that it plays a role in cell survival in this condition. This review enumerates and analyses the recent evidence that points to the role of ERR α as a regulator of hypoxic genes, both in cooperation with HIF-1 and through HIF-1-independent mechanisms, in invertebrate and vertebrate models and in physiological and pathological scenarios. ERR α 's functions during hypoxia include two mechanisms: (1) direct ERR α /HIF-1 interaction, which enhances HIF-1's transcriptional activity; and (2) transcriptional activation by the ERR α of genes that are classical HIF-1 targets, such as VEGF or glycolytic enzymes. ERR α is thus gaining recognition for its prominent role in the hypoxia response, both in the presence and absence of HIF-1. In some models, ERR α prepares cells for hypoxia, with important clinical/therapeutic implications.

Keywords: ERR; HIF-independent response to hypoxia; cancer; metabolic adaptation to hypoxia; VEGF; angiogenesis; ischemia; PGC-1 α

Citation: Chaltel-Lima, L.A.; Domínguez, F.; Domínguez-Ramírez, L.; Cortes-Hernandez, P. The Role of the Estrogen-Related Receptor alpha (ERR α) in Hypoxia and Its Implications for Cancer Metabolism. *Int. J. Mol. Sci.* **2023**, *24*, x. <https://doi.org/10.3390/xxxxx>

Academic Editor(s):

Received: date

Revised: date

Accepted: date

Published: date



Copyright: © 2023 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Oxygen, the final electron acceptor of the mitochondrial respiratory chain for ATP production, is crucial for all aerobic organisms. Despite the complex systems that higher organisms have developed to irrigate every organ and constantly provide all cells with oxygen, a variety of conditions can limit oxygen levels: some pathological (ischemia, tumor development, anemia, lung disease) and some physiological (embryonic development, exercise). Hypoxia is thus defined as a decrease in the oxygen supply to levels insufficient for cellular function [1]. The precise O₂ concentration that represents hypoxia varies from tissue to tissue, and likely even between individuals, as different tissues are exposed to different physiological oxygen concentrations (termed “physioxia”), most ranging from 3 to 9% oxygen (23–70 mmHg) [1]. O₂ determinations suggest that most mammalian high-energy-demand tissues, such as the brain, muscle, liver, renal cortex and heart, maintain physiological oxygen concentrations between 2.5 and 5.5% [1,2]. Since hypoxia can quickly become life threatening, it triggers a response to modulate blood flow, change energy metabolism, induce angiogenesis and cell differentiation and, ultimately, induce apoptosis [3].

The hypoxia response has a fast component that relies on existing proteins, such as ion channels and already-expressed signaling pathways, with effects such as blood

redistribution, tachypnea/tachycardia, widespread inhibition of protein translation and impaired cell proliferation [4,5]. It also has a well-characterized slower component that induces the expression of about one-thousand specific hypoxia genes once hypoxia is installed [6]. Hypoxia-inducible factors (HIFs) are central transcription factors responsible for protein expression during the slow component of the hypoxia response [7], but other transcription factors also participate. One such example is the estrogen-related receptor alpha (ERR α), a ubiquitously expressed orphan nuclear receptor, abundant in high-energy-demand tissues such as the heart, kidneys and cancer cells [8–11]. This review discusses the role that ERR α plays in the hypoxia response in synergism with HIF-1 and by HIF-independent mechanisms.

2. HIF-1, -2 and -3 Mediate the Hypoxia Response

The HIF-mediated transcriptional response to hypoxia was discovered in the 1990s [7] and received the Nobel Prize in Physiology or Medicine in 2019 [12]. Due to its central importance to physiology and to pathological states such as cancer, HIF has been extensively studied and reviewed [6,12–15]. HIF-1 was the first such factor described and remains the most characterized, but HIF-2 and HIF-3 have been described as well [14,16]. All are heterodimeric basic helix–loop–helix transcription factors consisting of subunits α and β . Each α subunit is O₂-regulated, as it is constantly targeted for destruction during normoxia via the Von Hippel Lindau protein and the E2-ligase/ubiquitin proteasome pathway [12–15,17]. In turn, the HIF-1 β subunit (initially known as the aryl hydrocarbon nuclear translocator ARNT, UniProt P27540) is constitutively expressed [12–15] and can heterodimerize with the different oxygen-sensitive α subunits (HIF-1 α , HIF-2 α or HIF-3 α) to create tissue-specific HIF-1, HIF-2 or HIF-3 transcription factors. While HIF-1 α is conserved from Parazoa to vertebrates and expressed in most cells [13,14], HIF-2 α and -3 α are only present in vertebrates and expressed tissue-specifically [12,14,16].

HIF-1 α (HIF1A, UniProt Q16665), the most characterized homolog, is an 826-residue protein that locates to the cytoplasm during normoxia where it is constantly destroyed [12–15,17]. Under hypoxic conditions or in the presence of iron chelating agents, it translocates to the nucleus and dimerizes with HIF-1 β to form the functional HIF-1 complex that activates gene transcription [12–15,17]. HIF-1 α DNA binding activity and stabilization have half maximal responses between 1.5 and 2% O₂ and maximal response at 0.5% O₂, determined in human cultured cells [2]. HIF-1 requires co-activators/transactivators such as CREB binding protein (CBP), p300 [18,19] and others [14], and the complex binds to hypoxia response elements (HREs) with the 5'-RCGTG-3' consensus [14,20].

Around one-thousand genes have been identified as HIF-1 targets [21] and they can be grouped into two functional categories: those that increase oxygen supply to tissues and those that decrease oxygen consumption by tissues [6,15]. In the first category, HIF elicits an increase in oxygen delivery to tissues by triggering erythropoiesis and angiogenesis through the expression of erythropoietin, the hormone that controls red cell production and blood O₂-carrying capacity [7], and VEGF (vascular endothelial growth factor), the main protein that stimulates new blood vessel formation [22]. In the second category (decreasing oxygen consumption) are many genes that modify energy metabolism [23,24]. For example, in hypoxia, oxidative phosphorylation (OXPHOS) is restricted due to the lack of O₂, and cells shift to anaerobic glycolysis through the increased expression of glycolytic enzymes by HIF-1 [23]. To compensate for the much lower ATP generation per glucose molecule through glycolysis than through OXPHOS, HIF-1 activates transcription of the SLC2A1 and SLC2A3 genes coding the glucose transporters GLUT1 and GLUT3 that increase glucose uptake [23]. Moreover, to inhibit the conversion of pyruvate to acetyl CoA, HIF-1 activates gene transcription to decrease pyruvate flux to the Krebs cycle and increase lactate production. Examples of activated genes are the PDK1 gene encoding PDH kinase, which phosphorylates and inactivates the catalytic subunit of pyruvate dehydrogenase (PDH) [24], and the LDHA gene encoding lactate dehydrogenase A, which

directly catalyzes the conversion of pyruvate to lactate [20]. In this way, the HIF-1 response to hypoxia is, in part, executed through metabolic adaptation.

To form HIF-2, HIF1- β heterodimerizes with HIF-2 α (EPAS1, UniProt Q99814; also called endothelial PAS domain protein 1, HIF-1 α -like factor (HLF), HIF-1 α related factor (HRF) and member of the PAS superfamily-1 (MOP-1)). HIF-2 α has similarities to HIF-1 α in terms of domain structure, O₂-dependent degradation, DNA sequence recognition (also binds to hypoxia response elements, HREs) and heterodimerization, yet exhibits different effects over gene expression mostly due to tissue-specific expression and kinetics [12,14,16]. The kinetics of HIF-1 α and HIF-2 α suggest that the former exerts a more rapid response at oxygen levels around 1–2%, whereas HIF-2 α action occurs after prolonged hypoxia [25]. In contrast to HIF-1 α that expresses ubiquitously, HIF-2 α only expresses in certain tissues such as embryonic and adult vascular endothelia, lung, placenta, heart, renal interstitial cells and liver [16,26]. HIF-2 α also has specific coactivators such as NF- κ B essential modulator, and Ets1 that do not interact with HIF-1 α . Genes strongly activated by HIF-2 are erythropoietin, VEGF receptor 2, insulin-like growth factor-binding protein-2 and plasminogen activator inhibitor-1 [26,27]. HIF-2 acts more effectively than HIF-1 on erythropoietin and iron metabolism genes, whereas VEGF and GLUT1 are similarly activated by HIF-1 and HIF-2, and glycolytic enzymes are more activated by HIF-1 [26,27]. Thus, within the hypoxia response that requires gene expression, HIF-1 constitutes a faster metabolic component; in turn, HIF-2 is more effective on erythropoiesis control once hypoxia persists.

In turn, HIF-3 α (HIF3A, UniProt Q9Y2N7) can be present in different splice variants that depend upon the tissue, some of which are proposed to have negative regulatory functions on the hypoxia response [16,28]. This seems to be the case for the short HIF-3 α variant that is also called inhibitory PAS domain protein (IPAS), which expresses in corneal epithelium and putatively prevents vascularization there [16,28]. Thus, HIF-3 α could have specific regulatory roles that will not be further reviewed here, but that have been discussed in [16,28].

3. Introduction to the Estrogen-Related Receptors (ERR)

The orchestration of metabolic adaptation central for hypoxia survival seems to involve other transcription factors that specialize in the control of energy metabolism, such as the estrogen-related receptor (ERR) subfamily of nuclear receptors. Here, we describe the subfamily, with a focus on ERR α , and then analyze the evidence and mechanisms that link ERR α to the regulation of hypoxic metabolism and angiogenesis.

The ERR subfamily belongs to group III of the nuclear receptor superfamily (orphan nuclear receptors) [8,29]. In humans and most vertebrates, it comprises three members: ERR α (NR3B1), ERR β (NR3B2) and ERR γ (NR3B3). However, only one ERR gene has been found in invertebrates such as Urochordates, *Drosophila melanogaster*, and in the mosquito *Anopheles gambiae*, but none seem to exist in *Caenorhabditis elegans* [30–32]. A search in the Inparanoid database (version 9) confirmed that no ERR homologs are identifiable in nematodes [33].

The general structure of ERRs is common to nuclear receptors, including four functional domains: N-terminal (NTD), DNA-binding (DBD), hinge, and a putative ligand-binding domain (LBD) [8,10,34] (Figure 1A). The DBD comprises two cysteine-rich zinc finger motifs, which are required for DNA binding and recognize the ERR response elements (ERREs), composed by the sequence TNAAGGTCA [35–37]. The three members of the ERR family (α , β and γ) bear high similarity, particularly in the DBD and LBD domains [38], but they have somewhat different functions and their expression is tissue-specific. ERR γ and ERR β bear more similarity with each other than with ERR α [38]. ERR α is the most abundant member of the family, expressed in most cells, and with higher levels in those with high energy demand, especially in cells that oxidize fatty acids [35–37], compatible with its role in the transcriptional control of energy metabolism.

ERR α (ESRRA, UniProt P11474) is a 423-residue protein and the first orphan nuclear receptor identified in a 1988 screen for genes related to estrogen receptor alpha (ER α) [39], just a few years before the identification of HIF. Unlike estrogen receptors, no endogenous ligand has been described for ERRs; thus, ERR α , the first orphan nuclear receptor identified, remains among the “non-adopted” orphans [40,41]. Recently, it was reported that an endogenous 19-nor steroid estradienolone, found in the urine of pregnant women, can bind and act as an inverse agonist to ERR α and ERR γ [42]. There is still little information to discern if this could be the long-sought endogenous ligand of the family, but it seems unlikely due to the plethora of crucial functions that have been described for the ERRs and which do not require a ligand (reviewed below).




Figure 1. A. ERR α 's structure and function. (A) The general domain topology of ERR α and other estrogen-related receptors (NR3) includes an amino-terminal domain (NTD), a DNA binding domain, a hinge, and a putative ligand-binding domain (LBD). (B,C) ERR α 's LBD has been crystalized in the absence (B) and presence of ligand. (C) The binding site is lined by residues F232, F286, F399 and F414 (PDBIDs 1XB7 and 2PJL) providing bulky side chains that fill the ligand-binding pocket in the absence of ligand (B). Moreover, without ligand, helix 11 (H11) lifts away from the binding site; helix 12 (H12) is perpendicular to H11 and residue F414 occludes the binding site (B). (D) ERR α can

bind DNA as a (i) monomer or (ii) dimer. ERR α 's transcriptional activity increases in complex with co-activators, for example PGC-1 α or β (iii), and through posttranslational modifications, such as phosphorylation (iv), mediated by HER2-EGF. ERR α activity increases in metabolic stress, cold, fasting/nutrient deprivation, exercise and hypoxia.

Due to ERR α 's sequence identity to ER α , particularly in the DBD and LBD (68% and 37% residue identity, respectively [39]), it was initially suggested that these two receptors share common targets, co-regulatory proteins and sites of action [43,44]. However, through the combination of computational biology; ERR α silencing; interaction with the co-activators such as PGC-1 α ; DNA binding assays; chromatin immunoprecipitation with sequencing (CHIP-seq); and reporter gene approaches, the differences between ERR α 's and ER α 's functions have become apparent [8,36,37,45]. ERR α regulates a different set of genes to ER α and is not involved in estrogen response. ERR α is mainly involved in the transcriptional regulation of metabolic pathways spanning carbohydrate, lipid and amino acid metabolism, importantly through the regulation of genes for mitochondrial biogenesis, oxidative phosphorylation (OXPHOS) and fatty acid oxidation [8,10,11,37]. The other members of the ERR family also control aspects of metabolism, although in specific tissues [8,29,32,38]. Overall, ERRs occupy the promoters of over 700 genes that encode mitochondrial proteins, regulating mitochondrial biogenesis [8].

Specifically, ERR α binds to the promoters of glycolysis and tricarboxylic acid cycle (TCA) genes, and to OXPHOS genes such as ATP synthase b (ATPsynb), cytochrome c (CYCS), COX4, GABPA and adenine nucleotide translocator 1 (ANT1) [45]. ChIP-seq studies performed in mouse or human, liver, kidney, macrophages or cancer cell lines confirmed that ERR α can bind to promoters for OXPHOS (*Sdh* and *Sucla2*), TCA (*Fasn*), glycolysis/gluconeogenesis and lipid metabolism genes (*Gpam* and *Elovl6*) [37,46–49]. Furthermore, ERR α activates the promoters of β -oxidation genes, such as ACADM (medium-chain acyl co-A dehydrogenase) and CPT1A (carnitine palmytoyl transferase 1A), as well as the promoters of glutamine transporters and enzymes for glutamine synthesis and catabolism [35,45]. In summary, ERR α 's function can be described as activating gene expression to adapt energy production to physiological or pathological stress. ERR α 's functions in cellular metabolism have been reviewed in [8]. In breast cancer, ERR α 's transcriptional activities mediate metabolic adaptations leading to treatment resistance [47]. The metabolic programs it controls make ERR α an ideal contributor to the hypoxia response and a potential pharmacological target.

The other members of the ERR subfamily also modulate metabolism with complementary and sometimes opposite functions to ERR α [8,29]. ERR β has emerged as important in maintaining multipotency [38]. In breast cancer, ERR α and ERR γ seem to play opposing roles as modulators of cell metabolism: ERR γ activates TCA and OXPHOS while ERR α redirects energy metabolism to glycolysis and lactate production [8]. This balance of control is likely part of the mechanism at the core of the Warburg effect in many tumors, along with HIF-1 [8,11], but it is far from a simple on/off switch. Rather, it is a dynamic balance under tight control that is highly cell- and context-specific, where both members of the ERR family, ERR α and ERR γ , activate metabolic pathways facilitating cell survival and adaptation to the changing environment.

To bind DNA and to modulate target genes, ERRs can act as monomers, homodimers or heterodimers [36,37] (Figure 1D), although in live cells mainly homo or heterodimers have been associated to function [37,38]. ERR transcriptional activity is increased by members of the steroid receptor co-activator (SRC) family [43,50,51] and by the peroxisome proliferator-activated receptor gamma co-activator-1 (PGC-1) α and β [45,50] (Figure 1D). Interactions with the cofactors are mediated by ERR's LBD, particularly by helices 11 and 12 via leucine rich motifs (H11 and H12 in Figure 1B,C), also referred to in the literature as ERR's AF2 domain for "activation function 2" [50,53,54]. In particular, ERR α 's functions on metabolism are more dependent on PGC-1 α [50]. ERR α and PGC-1 α influence each other's expression, and both orchestrate the transcription of energy metabolism genes

[37,54]. Recently, details on $ERR\alpha$'s transcription initiation mechanisms have been clarified. $PGC1\alpha$ was essential for p300 and mediator recruitment to activate transcription when $ERR\alpha$ acted on chromatin, whereas on naked DNA $ERR\alpha$ established direct contact with initiation factor TFIID, and $PGC1\alpha$ did not further increase transcription [50,51]. While $ERR\alpha$ depends on $PGC-1\alpha$ to transcribe metabolic genes, $ERR\beta$ and γ can function independently of $PGC-1\alpha$ in stem cells and muscle [50], and other cofactors that interact with their AF2 domains, such as NCOA, replace $PGC-1\alpha$ [50,51].

Unlike $ER\alpha$, the ERRs do not need ligands to interact with its co-activators and to bind DNA (they are constitutively active), probably because the putative ligand-binding pocket (LBP) is occupied by residue side chains in a conformation favored by binding to the cofactors. In the empty $ERR\alpha$ crystal structures, the binding pocket is mainly occupied by the bulky phenolic ring of Phe232 (XRD structures number this residue as 328; however, numbering according to UniProt is used here), which corresponds to a less bulky Ala350 in $ER\alpha$ [41]. Despite this apparent lack of a prominent ligand-binding pocket [41,53], synthetic compounds can inhibit $ERR\alpha$'s constitutive activity; thus, they are considered $ERR\alpha$'s inverse agonists (i.e., compounds with affinity and intrinsic activity on the protein). Among the first synthetic inverse agonists described for $ERR\alpha$ was XCT790 (reported in 2004), a thiadiazole acrylamide, which alters $ERR\alpha/PGC-1\alpha$ signaling and is inactive against the rest of the ERRs and $ER\alpha$ [55]. Later, "compound 1a" (cyclohexylmethyl-(1-p-tolyl-1H-indol-3-yl)-amine) and "compound 29" (4-(4-[(5R)-2,4-dioxo-1,3-thiazolidin-5-yl]methyl)-2-methoxyphenoxy)-3-(trifluoromethyl)benzotrile) were synthesized and have been crystallized in complex with $ERR\alpha$'s LBD [56,57]. An analysis of these $ERR\alpha$ structures with inhibitors revealed a significantly larger ligand-binding pocket than in the empty LBD, created by the rearrangement of amino acid residues F232, F286, F399 and F414 (328, 382, 495 and 510 in XRD 2PJL and 1XB7). F232 and F414 change conformation significantly when $ERR\alpha$ admits a ligand (Figure 1B vs. Figure 1C). In addition, these structures suggest that the presence of the inverse agonists disrupts the interaction between $ERR\alpha$ and $PGC-1\alpha$, through the displacement of $ERR\alpha$'s helix, to a position that interferes with co-activator recruitment [56,57].

4. Evidence of $ERR\alpha$'s Participation in the Hypoxia Response

Next, we review the evidence for $ERR\alpha$'s participation in the hypoxia response in models that span invertebrates and vertebrates, and physiological and pathological scenarios. These studies have led to the discovery of HIF-dependent and independent mechanisms, including some that transcend $ERR\alpha$'s central role as a metabolic coordinator during stress.

4.1. $ERR\alpha$ Induces VEGF Expression during Muscle Ischemia and Other Models

The work that first pointed to $ERR\alpha$'s role in hypoxia came from the study of angiogenesis/ischemia where VEGF, a classical HIF-1 target central to angiogenesis, was discovered to also be inducible by $ERR\alpha$ in skeletal muscle [58]. Arany et al. first detected that $ERR\alpha$'s co-activator, $PGC-1\alpha$, was induced by hypoxia in vitro in various cell types, and in vivo in muscle [58]. Using a skeletal muscle ischemia model, these authors showed that transgenic animals overexpressing $PGC-1\alpha$ had increased angiogenesis with VEGF expression. $PGC-1\alpha/ERR\alpha$, but not other transcription factors co-activated by $PGC-1\alpha$, were necessary to increase VEGF expression, through a mechanism that neither depended on HIF response elements (HREs) nor affected HIF-1 expression/stability [58]. Furthermore, conserved $ERR\alpha$ response elements (ERRES) were identified in the first intron of the VEGF gene and were recognized by $PGC-1\alpha/ERR\alpha$ [58]. $ERR\alpha$'s ability to induce VEGF expression and angiogenesis, as well as platelet-derived growth factor (PDGF) and Angiopoietin 2, has been confirmed in other models [59–64]. Some studies suggest that the effect does not require HIF-1 [58,59], while others suggest that HIF-1 can increase $ERR\alpha$ expression [61], and that, in turn, $ERR\alpha$ suppression can decrease HIF-1 α [60].

In skeletal muscle, the alternatively spliced truncated isoforms of PGC-1 α , NT-PGC-1 α and PGC-1 α 4, induced VEGF expression by ERR α without increasing mitochondrial biogenesis [64] (Figure 2B). These PGC1 α isoforms bind ERR α but not other transcription factors, such as NRF-1 and NRF-2 [53,64]. This suggests a mechanism by which the PGC-1 α /ERR α axis can operate in hypoxia without increasing mitochondria (Figure 2A), which would likely be impaired in respiration due to the limited O₂ to act as a terminal acceptor for OXPHOS. Additionally, other authors have suggested that PGC-1 α could amplify intracellular hypoxia by activating mitochondrial biogenesis/OXPHOS as a mechanism to consume all remaining intracellular oxygen [65], thus precipitating hypoxia responses and stabilizing HIF-1 α .




Figure 2. Metabolic adaptation via ERR α /PGC1 α in normoxia (A) vs. hypoxia (B). (A) In normoxic nutrient deprivation, ERR α /PGC1 α activate mitochondrial biogenesis. (B) In hypoxia, PGC-1 α 's truncated isoform NT-PGC-1 α binds ERR α but prevents the engagement of other transcription factors, limiting mitochondrial biogenesis and favoring the expression of angiogenesis genes such as VEGF. Oxygen consumption by mitochondria favors local hypoxia with concomitant HIF-1 α stabilization.

Other recent studies suggest that some $ERR\alpha$'s responses to hypoxia in the skeletal muscle are dependent on HIF-1. $ERR\alpha$ is expressed during hindlimb muscle ischemia. Transgenic mice overexpressing $ERR\alpha$ in the skeletal muscle have faster revascularization with more muscle capillaries and higher artery/arteriole density after ischemia [66,67]. $ERR\alpha$ overexpression was also induced in C2C12 myotubes by oxygen deprivation (culture in 95% nitrogen, 5% CO_2), hypoxia-mimetics such as dimethyl-oxaloylglycine (DMOG) or cobalt chloride ($CoCl_2$), or by nutrient deprivation [66]. Further in vitro experiments showed that $ERR\alpha$ regulates angiogenic gene expression through promoter recognition in C2C12 myotubes, and pointed out that $ERR\alpha$'s expression was HIF-1-dependent [66], for which the authors predicted 12 putative HIF1A::ARNT response elements in the $ERR\alpha$ gene promoter [66]. Altogether, these authors suggest that HIF is involved in the hypoxic induction of $ERR\alpha$ in the skeletal muscle through the transcriptional regulation of $ERR\alpha$ expression. However, $ERR\alpha$'s activity was not explored under HIF-1 depleting or activating conditions.

4.2. $ERR\alpha$ in Brain and Spinal Cord Hypoxia/Ischemia

Studies with astrocytes treated with CORM2, a CO-releasing compound that imitates ischemic brain injury, showed that $ERR\alpha$ / $PGG1\alpha$ can increase VEGF expression independent of HIF-1 (that is, even in HIF-1 α -deficient cells) [68]. The treatment induces Heme Oxygenase-1 (HO-1) expression and its metabolites (CO and bilirubin) and promotes Ca^{2+} influx through L-type Ca^{2+} channels producing $CaMKK\beta$ -mediated $AMPK\alpha$ activation [68]. $AMPK\alpha$ increases NAMPT expression and NAD^+ synthesis, which in turn increases SIRT activity. $PGC-1\alpha$ can be deacetylated by SIRT1 [68], and once deacetylated it interacts with $ERR\alpha$ to increase mitochondrial biogenesis and oxygen consumption [68]. With this model, the authors previously suggested that oxygen consumption aggravates intracellular hypoxia, allowing HIF-1 α stabilization that further increases $ERR\alpha$ / $PGG1\alpha$ expression [61]. Using ChIP assays, the authors proposed that HIF-1 can stimulate $ERR\alpha$'s transcription by binding to a putative HIF-1 response elements (+539 to +542, 5'-CGTG-3') within the promoter region of the $ERR\alpha$ gene [61]. HIF-1 α knockdown blocked $ERR\alpha$'s expression but not $PGG1\alpha$'s in that HO-1 inducing model. Therefore, it is likely that HO-1 can stimulate VEGF both via HIF-1 α dependent and independent mechanisms, the latter involving $PGC-1\alpha$ / $ERR\alpha$ and calcium regulation through the Ca^{2+} / $CaMKK$ / $AMPK$ pathway [68]. These authors also propose some reciprocal and dynamic coordination between HIF-1 α , and $PGC-1\alpha$ / $ERR\alpha$ for VEGF expression in astrocytic ischemia, involving mitochondrial biogenesis [61].

Other processes such as spinal cord injury (SCI) can manifest with ischemia, which aggravates secondary injury and neurological dysfunction [69–71]. Therefore, the vascular response is critical for SCI repair and includes HIF-1 α and VEGF expression. In an SCI rat model, $ERR\alpha$ inhibition with XCT790 decreased VEGF and angiopoietin-2 expression [72], which in turn decreased endothelial cell proliferation, vascular density and produced histopathological changes to the spinal cord, such as inflammatory cell infiltration, hemorrhage and vacuolation, and fewer normal neurons, suggesting that $ERR\alpha$ activity is essential for SCI repair, in part by favoring adequate re-vascularization via VEGF [72]. In this model, it has not been explored whether $ERR\alpha$'s effects require HIF-1.

In the microglial cell line, BV2, pharmacological $ERR\alpha$ inhibition (with XCT790) or activation (with pyrido [1,2- α]-pyrimidin-4-one) were explored in combination with $CoCl_2$ to mimic the hypoxia that accompanies SCI. Hypoxia induced HIF-1 α and autophagy. $ERR\alpha$'s effects were similar with/without hypoxia although more pronounced in hypoxia. During hypoxia, $ERR\alpha$ inhibition increased autophagy markers and increased IL-6, TNF- α and IL-10 mRNAs, but decreased FNDC5 (fibronectin type III domain containing protein 5) expression. In turn, $ERR\alpha$'s activation decreased p38 MAPK phosphorylation. The authors suggest that $ERR\alpha$ helps maintain homeostasis in microglia during hypoxia by down-modulating autophagy and inflammation [73].

4.3. *ERRα* in Hypobaric Hypoxia

On the other hand, in a non-pathological process such as exposure to high altitude, the expression of *ERRα* and *PGC-1α* are downregulated and the cell suffers mitochondrial dysfunction [74]. Treatment with dexamethasone maintains *ERRα* and *PGC-1α* levels similar to normoxia. This effect allows adaptability to hypobaric hypoxia through the expression of *ERRα* transcripts, *Fis1*, *Drp1* and *Mfn2*, which are mitochondrial dynamics proteins, and in turn increases OXPHOS [74]. This suggests that *ERRα*-mediated protection of mitochondrial bioenergetics is required for adaptation to hypobaric hypoxia.




Figure 3. Proposed interactions between HIF-1 and *ERRα* pathways in hypoxia. *ERRα*/*PGC-1α* express hypoxia genes acting with HIF-1 or independent from HIF-1. Ca^{2+} activates *CaMKKβ* and *AMPKα*, increasing *SIRT1*. *PGC-1α* is activated by *SIRT1* deacetylation, decreasing its destruction through the proteasome.

4.4. *ERRα*'s Role in Cancer-Related Hypoxia

In parallel, *ERRα* has been extensively studied in solid tumors where blood vessels frequently become limiting to irrigate the tumor mass, leading to hypoxia. Thus, cancer represents another model where extensive evidence points to *ERRα*'s contribution to the hypoxia response. Cancer cells in solid tumors use typical HIF-1 orchestrated mechanisms to survive hypoxia [75], impacting angiogenesis, cancer stem cell maintenance, metabolic reprogramming, epithelial–mesenchymal transition (EMT), invasion, metastasis and resistance to therapy (radiation and chemotherapy) [75–77].

Simultaneously, the overexpression of *ERRα* has been associated with tumor aggressiveness and poor prognosis [78–80]. In 2002, Ariazi et al. suggested *ERRα* as a biomarker of unfavorable clinical prognosis in breast cancer, due to increased *ERRα* mRNA levels in primary tumor cells against normal mammary epithelial cells. *ERRα*'s expression correlated with *Her2/ErbB2*, a tyrosine kinase receptor amplified in 15% to 25% of breast

cancers that also confers aggressiveness [78] and that increases $ERR\alpha$'s transcriptional activity via phosphorylation through MEK/MAPK and PI3K/Akt [80,81].

Subsequent immunohistochemical analyses, mRNA quantification and gene expression profiles in several solid tumors (breast, cervix, colon, endometrium, ovary and prostate) are in agreement with Ariazi et al. and relate $ERR\alpha$ overexpression to cancer aggressiveness, increased risk of recurrence and lower survival [9,77,82–86].

In breast and prostate cancer, $ERR\alpha$ has been found to interact directly with HIF-1 with two main effects: (1) HIF-1 stabilization; and (2) an increase in the HIF-dependent expression of hypoxic genes [86,87]. This evidence has led to the suggestion that the direct $ERR\alpha$ -HIF interaction is another important mechanism by which $ERR\alpha$ contributes to the hypoxic response. The physical interaction between HIF and $ERR\alpha$ has been explored using a series of $ERR\alpha$ truncation mutants covering the N terminus, DBD, and LBD in GST pull down assays. Ao et al. suggested that the $ERR\alpha$'s DBD is involved in HIF binding [87]. The immunoprecipitation of cellular lysates with anti- $ERR\alpha$, from MDA-MB-435 breast cancer cells treated with the iron chelator dipyrindyl (DP) to stabilize endogenous HIF-1 α , showed that all three ERRs associate to HIF α / β heterodimers both in vitro and in vivo, and this was abolished in $ERR\alpha$ mutants or with $ERR\alpha$ inhibitors [87]. Subsequent studies using co-immunoprecipitation and FRET in prostate cancer cells confirmed that the interaction happens and that it increases HIF-1's transcription but disagree on the $ERR\alpha$ domains involved [86]. Zou et al. suggest that the domain required for interaction with HIF-1 is the AF-2 region in $ERR\alpha$'s LBD [86]. These authors further suggest that $ERR\alpha$'s co-activator PGC1 α may be necessary for its interaction with HIF-1, as the AF-2 region in $ERR\alpha$'s LBD is important for co-activator recruitment. These effects are prevented in $ERR\alpha$ knockdowns or with $ERR\alpha$'s inverse agonist XCT790 [86,87]. Evidence in prostate cancer cells suggests that the $ERR\alpha$ /HIF-1 α interaction reduces the proteosomal degradation of HIF-1 α [86]. These authors suggest that $ERR\alpha$ overexpression stabilizes HIF-1 α and enhances HIF-1 transcriptional activity even under normoxia, with these effects amplified in hypoxia, resulting in a mechanism for the pre-adaptation to hypoxia [86].

In parallel, Stein et al. reported that $ERR\alpha$ regulates VEGF expression in breast cancer cell lines [62], similar to what was described in the previous section in angiogenesis models. A modified PGC1 α that only binds to $ERR\alpha$ was used to induce VEGF expression in MDA-MB231 and MCF7 breast cancer cells, and this effect was abolished with $ERR\alpha$ knockdowns. A main ERRE was located within the transcribed region of the VEGF gene [62]. The positive regulation of VEGF by $ERR\alpha$ has also been observed in human breast tumors and in murine models [63], supporting that VEGF is a direct transcriptional target of $ERR\alpha$ in cancer, as in other cell types (reviewed in previous section).

In summary, $ERR\alpha$ overexpression enhances the hypoxia response in solid tumor models. It is likely that $ERR\alpha$ functions as an aggressiveness factor in cancer because it prepares cancer cells to resist metabolic stress and hypoxia. $ERR\alpha$ has been observed as active in immunosuppressive and immunoresistant tumors [88]. Cancer models have pointed to HIF-dependent mechanisms such as the physical interaction between $ERR\alpha$ /HIF-1 α , as well as HIF-independent mechanisms such as VEGF modulation by $ERR\alpha$.

4.5. $ERR\alpha$ and Kidney Hypoxia

Organs with high energy demand such as the brain, heart and kidney have low tolerance to hypoxia and are good models to evaluate $ERR\alpha$'s effects. Physiological oxygen gradients across the renal cortex and medulla participate in the mechanisms to concentrate urine [89,90]. The healthy human kidney cortex presents around 50 mmHg of oxygen pressure, while the medulla has much lower oxygen pressures between 10 and 20 mmHg [89]. Keppner et al. recently evaluated the transcriptome during hypoxia (24 h at 0.2% O₂) of the cortical kidney murine cell line mCCD(c11) [90]. They found over 3000 differentially expressed genes, many related to aerobic metabolism and ATP production through

mitochondria, and the hypoxia response was mainly driven by HIF-1 and not HIF-2. Interestingly, they knocked down $ERR\alpha$ and identified a reduced expression of some genes that typically function in hypoxia, such as *Egln3* (an alpha-ketoglutarate dependent hydroxylase that controls cell proliferation and transcription upon hypoxia) and *Serpin1* (plasminogen activator inhibitor-1, involved in the control of blood clotting) [90]. Since this regulation happened without a change in HIF-1 α , the model suggests that $ERR\alpha$ controls the expression of specific genes important for the hypoxia response.

4.6. Hypoxia in the Invertebrate Fly Model

D. melanogaster is tolerant to oxygen starvation and can survive hypoxia for long periods of time. As in humans, the hypoxia response is importantly mediated by HIF (called *simA* in *D. melanogaster*); thus, *D. melanogaster* has been a study model for hypoxia [91] and represents a vertebrate model with a recognizable ERR. Li et al. showed that, in addition to HIF, the single ERR present in flies (called dERR) is necessary for the hypoxic response in *D. melanogaster*, since less than 25% of dERR mutant flies survived hypoxia [92]. Using single and double dERR and dHIF-1 α mutants, they described genes sets that are important for hypoxia response and detected a subset of 282 dERR-dependent transcripts that are HIF-independent and whose expression changed in hypoxia, such as *Pgi*, *Pfk*, *GAPDH2* and *LDH* [91]. This work suggests that the dERR has a prominent HIF-independent role in hypoxia adaptation, particularly via the upregulation of glycolytic enzymes.

Additionally, dERR was found to bind dHIF and participate in the HIF-mediated expression of its subset of genes [92]. The binding was shown by two hybrid screen and GST pull-downs and required dHIF's residues 1289–1293 (LKNLL) and dERR's LBD [92], in accordance with Zou et al. in cancer cells [86].

5. Conclusions

HIF-1 is the transcription factor usually considered the main regulator of the hypoxia response. However, $ERR\alpha$, a cellular metabolism regulator, also plays a key role in hypoxia survival, in models ranging from invertebrates to vertebrates and in physiological and pathological scenarios. $ERR\alpha$'s functions in hypoxia in most models include two mechanisms: (1) direct $ERR\alpha$ /HIF-1 interaction, which enhances HIF-1's transcriptional activity at HREs (possibly without $ERR\alpha$'s direct interaction with DNA); and (2) transcriptional activation by the $ERR\alpha$ of genes that are classical HIF-1 targets, such as VEGF or glycolytic enzymes. The second mechanism can even happen in a HIF-1 independent manner that depends on ERREs coexisting with HREs.

$ERR\alpha$ is thus gaining recognition for its prominent role in the hypoxia response, both in the presence and absence of HIF-1. In many models, $ERR\alpha$ prepares cells for hypoxia, with important clinical/therapeutical implications and perspectives that could allow for the manipulation of tissues so they are pre-adapted to resist hypoxia. This is important, as hypoxia is central to numerous diseases with significant human mortality and high costs, such as cancer, cardiovascular and pulmonary disease, stroke, bacterial infections, inflammation, disorders related to prematurity and wound healing.

ERR 's expression and activity are conserved from flies (and likely nematodes) to mammals, suggesting that the ERR-mediated response to hypoxia appeared early in evolution. Phylogenetic exploration of the ERR-HIF interaction warrants more interrogation, with the potential to yield insights into its mechanisms and how they evolved.

Despite the many models that have described $ERR\alpha$ as responding to ischemia, it is unknown if $ERR\alpha$ may directly sense oxygen. No mechanism for direct oxygen sensing by $ERR\alpha$ has been described. Alternatively, $ERR\alpha$'s activation upon ischemia/hypoxia may arise from the metabolic signals derived from ischemia or through HIF-1 stimulation. Protein tyrosine phosphatase 1B (PTP1B) and Parkin have also been shown to, respectively, decrease and increase the transcriptional activity of $ERR\alpha$ in hypoxia models

(pancreatic islets [93] and HeLa cells [94]); thus, other pathways and layers for ERR α modulation likely exist.

Author Contributions: L.A.C.-L.: Conceptualization, Writing—original draft preparation; F.D.: Resources, Writing—reviewing and validation; L.D.-R.: Conceptualization, Software, Writing—reviewing and editing; P.C.-H.: Supervision, Conceptualization, Writing—reviewing and editing. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by IMSS, grant number FIS/IMSS/PROT/G16/1591, and CONACYT FOSISS 261918. LACL received research scholarships from CONACYT and IMSS.

Institutional Review Board Statement:

Informed Consent Statement:

Data Availability Statement:

Acknowledgments: LDR expresses his gratitude to Dra. Marietta Tuena Sangri, the most wonderful mentor any young scientist could ever hope to have. She will be dearly missed.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Carreau, A.; Hafny-Rahbi, B.E.; Matejuk, A.; Grillon, C.; Kieda, C. Why Is the Partial Oxygen Pressure of Human Tissues a Crucial Parameter? Small Molecules and Hypoxia. *J. Cell Mol. Med.* **2011**, *15*, 1239–1253. <https://doi.org/10.1111/j.1582-4934.2011.01258.x>.
2. Jiang, B.-H.; Zheng, J.Z.; Leung, S.W.; Roe, R.; Semenza, G.L. Transactivation and Inhibitory Domains of Hypoxia-Inducible Factor 1 α Modulation of Transcriptional Activity by Oxygen Tension*. *J. Biol. Chem.* **1997**, *272*, 19253–19260. <https://doi.org/10.1074/jbc.272.31.19253>.
3. Rocha, S. Gene Regulation under Low Oxygen: Holding Your Breath for Transcription. *Trends Biochem. Sci.* **2007**, *32*, 389–397. <https://doi.org/10.1016/j.tibs.2007.06.005>.
4. Seta, K.A.; Spicer, Z.; Yuan, Y.; Lu, G.; Millhorn, D.E. Responding to Hypoxia: Lessons from a Model Cell Line. *Sci. Stke.* **2002**, *2002*, re11. <https://doi.org/10.1126/stke.2002.146.re11>.
5. Fels, D.R.; Koumenis, C. The PERK/eIF2 α /ATF4 Module of the UPR in Hypoxia Resistance and Tumor Growth. *Cancer Biol. Ther.* **2006**, *5*, 723–728. <https://doi.org/10.4161/cbt.5.7.2967>.
6. Semenza, G.L. Hypoxia-Inducible Factors in Physiology and Medicine. *Cell* **2012**, *148*, 399–408. <https://doi.org/10.1016/j.cell.2012.01.021>.
7. Semenza, G.L.; Wang, G.L. A Nuclear Factor Induced by Hypoxia via de Novo Protein Synthesis Binds to the Human Erythropoietin Gene Enhancer at a Site Required for Transcriptional Activation. *Mol. Cell Biol.* **1992**, *12*, 5447–5454. <https://doi.org/10.1128/mcb.12.12.5447>.
8. Tam, I.S.; Giguère, V. There and Back Again: The Journey of the Estrogen-Related Receptors in the Cancer Realm. *J. Steroid Biochem. Mol. Biol.* **2016**, *157*, 13–19. <https://doi.org/10.1016/j.jsbmb.2015.06.009>.
9. Matsushima, H.; Mori, T.; Ito, F.; Yamamoto, T.; Akiyama, M.; Kokabu, T.; Yoriki, K.; Umemura, S.; Akashi, K.; Kitawaki, J. Anti-Tumor Effect of Estrogen-Related Receptor Alpha Knockdown on Uterine Endometrial Cancer. *Oncotarget* **2016**, *7*, 34131–34148. <https://doi.org/10.18632/oncotarget.9151>.
10. Deblois, G.; Giguère, V. Oestrogen-Related Receptors in Breast Cancer: Control of Cellular Metabolism and Beyond. *Nat. Rev. Cancer* **2013**, *13*, 27–36. <https://doi.org/10.1038/nrc3396>.
11. Cai, Q.; Lin, T.; Kamarajugadda, S.; Lu, J. Regulation of Glycolysis and the Warburg Effect by Estrogen-Related Receptors. *Oncogene* **2013**, *32*, 2079–2086. <https://doi.org/10.1038/onc.2012.221>.
12. Advanced Information. The Nobel Prize in Physiology or Medicine 2019. NobelPrize.org Nobel Prize Outreach AB. 2023. Available online: <https://www.nobelprize.org/prizes/medicine/2019/advanced-information/> (accessed on).
13. Semenza, G.L. Oxygen Sensing, Homeostasis, and Disease. *N. Engl. J. Med.* **2011**, *365*, 537–547. <https://doi.org/10.1056/nejmra1011165>.
14. Semenza, G.L. The Genomics and Genetics of Oxygen Homeostasis. *Annu. Rev. Genom. Hum. Genet.* **2020**, *21*, 183–204. <https://doi.org/10.1146/annurev-genom-111119-073356>.
15. Soni, S.; Padwad, Y.S. HIF-1 in Cancer Therapy: Two Decade Long Story of a Transcription Factor. *Acta Oncol.* **2017**, *56*, 503–515. <https://doi.org/10.1080/0284186x.2017.1301680>.
16. Albadari, N.; Deng, S.; Li, W. The Transcriptional Factors HIF-1 and HIF-2 and Their Novel Inhibitors in Cancer Therapy. *Expert. Opin. Drug Dis.* **2019**, *14*, 667–682. <https://doi.org/10.1080/17460441.2019.1613370>.
17. Chowdhury, R.; Leung, I.K.H.; Tian, Y.-M.; Abboud, M.I.; Ge, W.; Domene, C.; Cantrelle, F.-X.; Landrieu, I.; Hardy, A.P.; Pugh, C.W.; et al. Structural Basis for Oxygen Degradation Domain Selectivity of the HIF Prolyl Hydroxylases. *Nat. Commun.* **2016**, *7*, 12673. <https://doi.org/10.1038/ncomms12673>.

18. Kallio, P.J.; Okamoto, K.; O'Brien, S.; Carrero, P.; Makino, Y.; Tanaka, H.; Poellinger, L. Signal Transduction in Hypoxic Cells: Inducible Nuclear Translocation and Recruitment of theCBP/P300 Coactivator by the Hypoxia-induciblefactor-1 α . *EMBO J.* **1998**, *17*, 6573–6586. <https://doi.org/10.1093/emboj/17.22.6573>.
19. Bhattacharya, S.; Michels, C.L.; Leung, M.-K.; Arany, Z.P.; Kung, A.L.; Livingston, D.M. Functional Role of P35srj, a Novel P300/CBP Binding Protein, during Transactivation by HIF-1. *Gene Dev.* **1999**, *13*, 64–75. <https://doi.org/10.1101/gad.13.1.64>.
20. Semenza, G.L.; Jiang, B.-H.; Leung, S.W.; Passantino, R.; Concordet, J.-P.; Maire, P.; Giallongo, A. Hypoxia Response Elements in the Aldolase A, Enolase 1, and Lactate Dehydrogenase A Gene Promoters Contain Essential Binding Sites for Hypoxia-Inducible Factor 1*. *J. Biol. Chem.* **1996**, *271*, 32529–32537. <https://doi.org/10.1074/jbc.271.51.32529>.
21. Xia, X.; Lemieux, M.E.; Li, W.; Carroll, J.S.; Brown, M.; Liu, X.S.; Kung, A.L. Integrative Analysis of HIF Binding and Transactivation Reveals Its Role in Maintaining Histone Methylation Homeostasis. *Proc. Natl. Acad. Sci. USA* **2009**, *106*, 4260–4265. <https://doi.org/10.1073/pnas.0810067106>.
22. Forsythe, J.A.; Jiang, B.-H.; Iyer, N.V.; Agani, F.; Leung, S.W.; Koos, R.D.; Semenza, G.L. Activation of Vascular Endothelial Growth Factor Gene Transcription by Hypoxia-Inducible Factor 1. *Mol. Cell. Biol.* **1996**, *16*, 4604–4613. <https://doi.org/10.1128/mcb.16.9.4604>.
23. Iyer, N.V.; Kotch, L.E.; Agani, F.; Leung, S.W.; Laughner, E.; Wenger, R.H.; Gassmann, M.; Gearhart, J.D.; Lawler, A.M.; Yu, A.Y.; et al. Cellular and Developmental Control of O₂ Homeostasis by Hypoxia-Inducible Factor 1 α . *Gene Dev.* **1998**, *12*, 149–162. <https://doi.org/10.1101/gad.12.2.149>.
24. Papandreou, I.; Cairns, R.A.; Fontana, L.; Lim, A.L.; Denko, N.C. HIF-1 Mediates Adaptation to Hypoxia by Actively Downregulating Mitochondrial Oxygen Consumption. *Cell Metab.* **2006**, *3*, 187–197. <https://doi.org/10.1016/j.cmet.2006.01.012>.
25. Pählman, S.; Mohlin, S. Hypoxia and Hypoxia-Inducible Factors in Neuroblastoma. *Cell Tissue Res.* **2018**, *372*, 269–275. <https://doi.org/10.1007/s00441-017-2701-1>.
26. Haase, V.H. Regulation of Erythropoiesis by Hypoxia-Inducible Factors. *Blood Rev.* **2013**, *27*, 41–53. <https://doi.org/10.1016/j.blre.2012.12.003>.
27. Loboda, A.; Jozkowicz, A.; Dulak, J. HIF-1 and HIF-2 Transcription Factors—Similar but Not Identical. *Mol. Cells* **2010**, *29*, 435–442. <https://doi.org/10.1007/s10059-010-0067-2>.
28. Duan, C. Hypoxia-Inducible Factor 3 Biology: Complexities and Emerging Themes. *Am. J. Physiol.-Cell Physiol.* **2016**, *310*, C260–C269. <https://doi.org/10.1152/ajpcell.00315.2015>.
29. Bookout, A.L.; Jeong, Y.; Downes, M.; Yu, R.T.; Evans, R.M.; Mangelsdorf, D.J. Anatomical Profiling of Nuclear Receptor Expression Reveals a Hierarchical Transcriptional Network. *Cell* **2006**, *126*, 789–799. <https://doi.org/10.1016/j.cell.2006.06.049>.
30. Park, W.; Kim, G.J.; Choi, H.; Vanacker, J.-M.; Sohn, Y.C. Conserved Properties of a Urochordate Estrogen Receptor-Related Receptor (ERR) with Mammalian ERR α . *Biochim. Biophys. Acta BBA—Gene Regul. Mech.* **2009**, *1789*, 125–134. <https://doi.org/10.1016/j.bbagr.2008.08.011>.
31. Bardet, P.-L.; Laudet, V.; Vanacker, J.-M. Studying Non-Mammalian Models? Not a Fool's ERRand! *Trends Endocrinol. Metab.* **2006**, *17*, 166–171. <https://doi.org/10.1016/j.tem.2006.03.005>.
32. Maglich, J.M.; Sluder, A.; Guan, X.; Shi, Y.; McKee, D.D.; Carrick, K.; Kamdar, K.; Willson, T.M.; Moore, J.T. Comparison of Complete Nuclear Receptor Sets from the Human, Caenorhabditis Elegans and Drosophila Genomes. *Genome Biol.* **2001**, *2*, research0029.1. <https://doi.org/10.1186/gb-2001-2-8-research0029>.
33. Persson, E.; Sonnhammer, E.L.L. InParanoidDB 9: Ortholog Groups for Protein Domains and Full-Length Proteins. *J. Mol. Biol.* **2023**, 168001. <https://doi.org/10.1016/j.jmb.2023.168001>.
34. Kallen, J.; Lattmann, R.; Beerli, R.; Blechschmidt, A.; Blommers, M.J.J.; Geiser, M.; Ottl, J.; Schlaeppli, J.-M.; Strauss, A.; Fournier, B. Crystal Structure of Human Estrogen-Related Receptor Alpha in Complex with a Synthetic Inverse Agonist Reveals Its Novel Molecular Mechanism. *J. Biol. Chem.* **2007**, *282*, 23231–23239. <https://doi.org/10.1074/jbc.m703337200>.
35. Sladek, R.; Bader, J.A.; Giguère, V. The Orphan Nuclear Receptor Estrogen-Related Receptor Alpha Is a Transcriptional Regulator of the Human Medium-Chain Acyl Coenzyme A Dehydrogenase Gene. *Mol. Cell Biol.* **1997**, *17*, 5400–5409. <https://doi.org/10.1128/mcb.17.9.5400>.
36. Barry, J.B.; Laganière, J.; Giguère, V. A Single Nucleotide in an Estrogen-Related Receptor α Site Can Dictate Mode of Binding and Peroxisome Proliferator-Activated Receptor γ Coactivator 1 α Activation of Target Promoters. *Mol. Endocrinol.* **2006**, *20*, 302–310. <https://doi.org/10.1210/me.2005-0313>.
37. Dufour, C.R.; Wilson, B.J.; Huss, J.M.; Kelly, D.P.; Alaynick, W.A.; Downes, M.; Evans, R.M.; Blanchette, M.; Giguère, V. Genome-Wide Orchestration of Cardiac Functions by the Orphan Nuclear Receptors ERR α and γ . *Cell Metab.* **2007**, *5*, 345–356. <https://doi.org/10.1016/j.cmet.2007.03.007>.
38. Festuccia, N.; Owens, N.; Navarro, P. Esrrb, an Estrogen-Related Receptor Involved in Early Development, Pluripotency, and Reprogramming. *FEBS Lett.* **2018**, *592*, 852–877. <https://doi.org/10.1002/1873-3468.12826>.
39. Giguère, V.; Yang, N.; Segui, P.; Evans, R.M. Identification of a New Class of Steroid Hormone Receptors. *Nature* **1988**, *331*, 91–94. <https://doi.org/10.1038/331091a0>.
40. Ingraham, H.A.; Redinbo, M.R. Orphan Nuclear Receptors Adopted by Crystallography. *Curr. Opin. Struc. Biol.* **2005**, *15*, 708–715. <https://doi.org/10.1016/j.sbi.2005.10.009>.
41. Kallen, J.; Schlaeppli, J.-M.; Bitsch, F.; Filipuzzi, I.; Schilb, A.; Riou, V.; Graham, A.; Strauss, A.; Geiser, M.; Fournier, B. Evidence for Ligand-Independent Transcriptional Activation of the Human Estrogen-Related Receptor Alpha (ERR α): Crystal

- Structure of ERR α Ligand Binding Domain in Complex with Peroxisome Proliferator-Activated Receptor Coactivator-1 α . *J. Biol. Chem.* **2004**, *279*, 49330–49337. <https://doi.org/10.1074/jbc.m407999200>.
42. Ghanbari, F.; Hebert-Losier, A.; Barry, J.; Poirier, D.; Giguere, V.; Mader, S.; Philip, A. Isolation and Functional Characterization of a Novel Endogenous Inverse Agonist of Estrogen Related Receptors (ERRs) from Human Pregnancy Urine. *J. Steroid. Biochem. Mol. Biol.* **2019**, *191*, 105352. <https://doi.org/10.1016/j.jsbmb.2019.04.001>.
43. Zhang, Z.; Teng, C.T. Estrogen Receptor-Related Receptor A1 Interacts with Coactivator and Constitutively Activates the Estrogen Response Elements of the Human Lactoferrin Gene*. *J. Biol. Chem.* **2000**, *275*, 20837–20846. <https://doi.org/10.1074/jbc.m001880200>.
44. Giguère, V. To ERR in the Estrogen Pathway. *Trends Endocrinol. Metab.* **2002**, *13*, 220–225. [https://doi.org/10.1016/s1043-2760\(02\)00592-1](https://doi.org/10.1016/s1043-2760(02)00592-1).
45. Gaillard, S.; Grasfeder, L.L.; Haeffele, C.L.; Lobenhofer, E.K.; Chu, T.-M.; Wolfinger, R.; Kazmin, D.; Koves, T.R.; Muoio, D.M.; Chang, C.; et al. Receptor-Selective Coactivators as Tools to Define the Biology of Specific Receptor-Coactivator Pairs. *Mol. Cell* **2006**, *24*, 797–803. <https://doi.org/10.1016/j.molcel.2006.10.012>.
46. Chaveroux, C.; Eichner, L.J.; Dufour, C.R.; Shatnawi, A.; Khoutorsky, A.; Bourque, G.; Sonenberg, N.; Giguère, V. Molecular and Genetic Crosstalks between mTOR and ERR α Are Key Determinants of Rapamycin-Induced Nonalcoholic Fatty Liver. *Cell Metab.* **2013**, *17*, 586–598. <https://doi.org/10.1016/j.cmet.2013.03.003>.
47. Deblois, G.; Hall, J.A.; Perry, M.-C.; Laganière, J.; Ghahremani, M.; Park, M.; Hallett, M.; Giguère, V. Genome-Wide Identification of Direct Target Genes Implicates Estrogen-Related Receptor Alpha as a Determinant of Breast Cancer Heterogeneity. *Cancer Res.* **2009**, *69*, 6149–6157. <https://doi.org/10.1158/0008-5472.can-09-1251>.
48. Tremblay, A.M.; Dufour, C.R.; Ghahremani, M.; Reudelhuber, T.L.; Giguère, V. Physiological Genomics Identifies Estrogen-Related Receptor α as a Regulator of Renal Sodium and Potassium Homeostasis and the Renin-Angiotensin Pathway. *Mol. Endocrinol.* **2010**, *24*, 22–32. <https://doi.org/10.1210/me.2009-0254>.
49. Deblois, G.; Smith, H.W.; Tam, I.S.; Gravel, S.-P.; Caron, M.; Savage, P.; Labbé, D.P.; Bégin, L.R.; Tremblay, M.L.; Park, M.; et al. ERR α Mediates Metabolic Adaptations Driving Lapatinib Resistance in Breast Cancer. *Nat. Commun.* **2016**, *7*, 12156. <https://doi.org/10.1038/ncomms12156>.
50. Giguère, V. Transcription Initiation by the ERRs: No Ligand but Two Activation Pathways. *Cell Res.* **2023**, *33*, 269–270. <https://doi.org/10.1038/s41422-023-00780-9>.
51. Nakadai, T.; Shimada, M.; Ito, K.; Cevher, M.A.; Chu, C.-S.; Kumegawa, K.; Maruyama, R.; Malik, S.; Roeder, R.G. Two Target Gene Activation Pathways for Orphan ERR Nuclear Receptors. *Cell Res.* **2023**, *33*, 165–183. <https://doi.org/10.1038/s41422-022-00774-z>.
52. Sonoda, J.; Laganière, J.; Mehl, I.R.; Barish, G.D.; Chong, L.-W.; Li, X.; Scheffler, I.E.; Mock, D.C.; Bataille, A.R.; Robert, F.; et al. Nuclear Receptor ERR α and Coactivator PGC-1 β Are Effectors of IFN- γ -Induced Host Defense. *Gene Dev.* **2007**, *21*, 1909–1920. <https://doi.org/10.1101/gad.1553007>.
53. Takacs, M.; Petoukhov, M.V.; Atkinson, R.A.; Roblin, P.; Ogi, F.-X.; Demeler, B.; Potier, N.; Chebaro, Y.; Dejaegere, A.; Svergun, D.I.; et al. The Asymmetric Binding of PGC-1 α to the ERR α and ERR γ Nuclear Receptor Homodimers Involves a Similar Recognition Mechanism. *PLoS ONE* **2013**, *8*, e67810. <https://doi.org/10.1371/journal.pone.0067810>.
54. Sihag, S.; Cresci, S.; Li, A.Y.; Sucharov, C.C.; Lehman, J.J. PGC-1 α and ERR α Target Gene Downregulation Is a Signature of the Failing Human Heart. *J. Mol. Cell Cardiol.* **2009**, *46*, 201–212. <https://doi.org/10.1016/j.yjmcc.2008.10.025>.
55. Busch, B.B.; Stevens, W.C.; Martin, R.; Ordentlich, P.; Zhou, S.; Sapp, D.W.; Horlick, R.A.; Mohan, R. Identification of a Selective Inverse Agonist for the Orphan Nuclear Receptor Estrogen-Related Receptor Alpha. *J. Med. Chem.* **2004**, *47*, 5593–5596. <https://doi.org/10.1021/jm049334f>.
56. Kallen, J.; Lattmann, R.; Beerli, R.; Blechschmidt, A.; Blommers, M.J.J.; Geiser, M.; Ottl, J.; Schlaeppli, J.-M.; Strauss, A.; Fournier, B. Crystal Structure of Human Estrogen-Related Receptor Alpha in Complex with a Synthetic Inverse Agonist Reveals Its Novel Molecular Mechanism. *J. Biol. Chem.* **2007**, *282*, 23231–23239. <https://doi.org/10.1074/jbc.m703337200>.
57. Patch, R.J.; Searle, L.L.; Kim, A.J.; De, D.; Zhu, X.; Askari, H.B.; O'Neill, J.C.; Abad, M.C.; Rentzeperis, D.; Liu, J.; et al. Identification of Diaryl Ether-Based Ligands for Estrogen-Related Receptor α as Potential Antidiabetic Agents. *J. Med. Chem.* **2011**, *54*, 788–808. <https://doi.org/10.1021/jm101063h>.
58. Arany, Z.; Foo, S.-Y.; Ma, Y.; Ruas, J.L.; Bommi-Reddy, A.; Girnun, G.; Cooper, M.; Laznik, D.; Chinsomboon, J.; Rangwala, S.M.; et al. HIF-Independent Regulation of VEGF and Angiogenesis by the Transcriptional Coactivator PGC-1 α . *Nature* **2008**, *451*, 1008–1012. <https://doi.org/10.1038/nature06613>.
59. Zhang, K.; Lu, J.; Mori, T.; Smith-Powell, L.; Synold, T.W.; Chen, S.; Wen, W. Baicalin Increases VEGF Expression and Angiogenesis by Activating the ERR α /PGC-1 α Pathway. *Cardiovasc. Res.* **2011**, *89*, 426–435. <https://doi.org/10.1093/cvr/cvq296>.
60. Zhang, L.-D.; Chen, L.; Zhang, M.; Qi, H.-J.; Chen, L.; Chen, H.-F.; Zhong, M.-K.; Shi, X.-J.; Li, Q.-Y. Downregulation of ERR α Inhibits Angiogenesis in Human Umbilical Vein Endothelial Cells through Regulating VEGF Production and PI3K/Akt/STAT3 Signaling Pathway. *Eur. J. Pharmacol.* **2015**, *769*, 167–176. <https://doi.org/10.1016/j.ejphar.2015.11.014>.
61. Choi, Y.K.; Park, J.H.; Yun, J.-A.; Cha, J.-H.; Kim, Y.; Won, M.-H.; Kim, K.-W.; Ha, K.-S.; Kwon, Y.-G.; Kim, Y.-M. Heme Oxygenase Metabolites Improve Astrocytic Mitochondrial Function via a Ca²⁺-Dependent HIF-1 α /ERR α Circuit. *PLoS ONE* **2018**, *13*, e0202039. <https://doi.org/10.1371/journal.pone.0202039>.
62. Stein, R.A.; Gaillard, S.; McDonnell, D.P. Estrogen-Related Receptor Alpha Induces the Expression of Vascular Endothelial Growth Factor in Breast Cancer Cells. *J. Steroid Biochem. Mol. Biol.* **2009**, *114*, 106–112. <https://doi.org/10.1016/j.jsbmb.2009.02.010>.

63. Fradet, A.; Sorel, H.; Bouazza, L.; Goehrig, D.; Dépalle, B.; Bellahcène, A.; Castronovo, V.; Follet, H.; Descotes, F.; Aubin, J.E.; et al. Dual Function of $ERR\alpha$ in Breast Cancer and Bone Metastasis Formation: Implication of VEGF and Osteoprotegerin. *Cancer Res.* **2011**, *71*, 5728–5738. <https://doi.org/10.1158/0008-5472.can-11-1431>.
64. Thom, R.; Rowe, G.C.; Jang, C.; Safdar, A.; Arany, Z. Hypoxic Induction of Vascular Endothelial Growth Factor (VEGF) and Angiogenesis in Muscle by Truncated Peroxisome Proliferator-Activated Receptor γ Coactivator (PGC)-1 α . *J. Biol. Chem.* **2014**, *289*, 8810–8817. <https://doi.org/10.1074/jbc.m114.554394>.
65. Sancho, P.; Burgos-Ramos, E.; Tavera, A.; Bou Kheir, T.; Jagust, P.; Schoenhals, M.; Barneda, D.; Sellers, K.; Campos-Olivas, R.; Graña, O.; et al. MYC/PGC-1 α Balance Determines the Metabolic Phenotype and Plasticity of Pancreatic Cancer Stem Cells. *Cell Metab.* **2015**, *22*, 590–605. <https://doi.org/10.1016/j.cmet.2015.08.015>.
66. Sopariwala, D.H.; Likhite, N.; Pei, G.; Haroon, F.; Lin, L.; Yadav, V.; Zhao, Z.; Narkar, V.A. Estrogen-related Receptor α Is Involved in Angiogenesis and Skeletal Muscle Revascularization in Hindlimb Ischemia. *FASEB J.* **2021**, *35*, e21480. <https://doi.org/10.1096/fj.202001794rr>.
67. Sopariwala, D.H.; Rios, A.S.; Park, M.K.; Song, M.S.; Kumar, A.; Narkar, V.A. Estrogen-related Receptor Alpha Is an AMPK-regulated Factor that Promotes Ischemic Muscle Revascularization and Recovery in Diet-induced Obese Mice. *FASEB Bioadv.* **2022**, *4*, 602–618. <https://doi.org/10.1096/fba.2022-00015>.
68. Choi, Y.K.; Kim, J.-H.; Lee, D.-K.; Lee, K.-S.; Won, M.-H.; Jeoung, D.; Lee, H.; Ha, K.-S.; Kwon, Y.-G.; Kim, Y.-M. Carbon Monoxide Potentiation of L-Type Ca^{2+} Channel Activity Increases HIF-1 α -Independent VEGF Expression via an AMPK α /SIRT1-Mediated PGC-1 α /ERR α Axis. *Antioxid. Redox Sign* **2017**, *27*, 21–36. <https://doi.org/10.1089/ars.2016.6684>.
69. Xiaowei, H.; Ninghui, Z.; Wei, X.; Yiping, T.; Linfeng, X. The Experimental Study of Hypoxia-Inducible Factor-1 α and Its Target Genes in Spinal Cord Injury. *Spinal Cord* **2006**, *44*, 35–43. <https://doi.org/10.1038/sj.sc.3101813.a56>.
70. Chen, M.-H.; Ren, Q.-X.; Yang, W.-F.; Chen, X.-L.; Lu, C.; Sun, J. Influences of HIF-1 α on Bax/Bcl-2 and VEGF Expressions in Rats with Spinal Cord Injury. *Int. J. Clin. Exp. Pathol.* **2013**, *6*, 2312–2322.a57.
71. Mortazavi, M.M.; Verma, K.; Harmon, O.A.; Griessenauer, C.J.; Adeeb, N.; Theodore, N.; Tubbs, R.S. The Microanatomy of Spinal Cord Injury: A Review. *Clin. Anat.* **2015**, *28*, 27–36. <https://doi.org/10.1002/ca.22432.a58>.
72. Hu, J.Z.; Long, H.; Wu, T.-D.; Zhou, Y.; Lu, H.-B. The Effect of Estrogen-Related Receptor α on the Regulation of Angiogenesis after Spinal Cord Injury. *Neuroscience* **2015**, *290*, 570–580. <https://doi.org/10.1016/j.neuroscience.2015.01.067>.
73. Deng, C.-Y.; Zhu, T.-T.; Lian, S.; Wang, J.-F.; Wu, R.; Zheng, J.-S. Estrogen-Related Receptor α (ERR α) Functions in the Hypoxic Injury of Microglial Cells. *J. Vet. Res.* **2022**, *66*, 131–140. <https://doi.org/10.2478/jvetres-2022-0009>.
74. Chitra, L.; Boopathy, R. Adaptability to Hypobaric Hypoxia Is Facilitated through Mitochondrial Bioenergetics: An In Vivo Study. *Brit. J. Pharmacol.* **2013**, *169*, 1035–1047. <https://doi.org/10.1111/bph.12179>.
75. Semenza, G.L. Defining the Role of Hypoxia-Inducible Factor 1 in Cancer Biology and Therapeutics. *Oncogene* **2010**, *29*, 625–634. <https://doi.org/10.1038/onc.2009.441>.
76. Wang, M.; Yang, G.; Jiang, X.; Lu, D.; Mei, H.; Chen, B. Peroxisome Proliferator-Activated Receptor- γ Coactivator-1 α (PGC-1 α) Regulates the Expression of B-Cell Lymphoma/Leukemia-2 (Bcl-2) and Promotes the Survival of Mesenchymal Stem Cells (MSCs) via PGC-1 α /ERR α Interaction in the Absence of Serum, Hypoxia, and High Glucose Conditions. *Med. Sci. Monit. Int. Med. J. Exp. Clin. Res.* **2017**, *23*, 3451–3460. <https://doi.org/10.12659/msm.902183>.
77. Zhong, H.; Marzo, A.M.D.; Laughner, E.; Lim, M.; Hilton, D.A.; Zagzag, D.; Buechler, P.; Isaacs, W.B.; Semenza, G.L.; Simons, J.W. Overexpression of Hypoxia-Inducible Factor 1 α in Common Human Cancers and Their Metastases. *Cancer Res.* **1999**, *59*, 5830–5835.
78. Ariazi, E.A.; Clark, G.M.; Mertz, J.E. Estrogen-Related Receptor Alpha and Estrogen-Related Receptor Gamma Associate with Unfavorable and Favorable Biomarkers, Respectively, in Human Breast Cancer. *Cancer Res.* **2002**, *62*, 6510–6518.
79. Suzuki, T.; Miki, Y.; Moriya, T.; Shimada, N.; Ishida, T.; Hirakawa, H.; Ohuchi, N.; Sasano, H. Estrogen-Related Receptor α in Human Breast Carcinoma as a Potent Prognostic Factor. *Cancer Res.* **2004**, *64*, 4670–4676. <https://doi.org/10.1158/0008-5472.can-04-0250>.
80. Ariazi, E.A.; Kraus, R.J.; Farrell, M.L.; Jordan, V.C.; Mertz, J.E. Estrogen-Related Receptor A1 Transcriptional Activities Are Regulated in Part via the ErbB2/HER2 Signaling Pathway. *Am. Assoc. Cancer Res* **2007**, *5*, 71–85. <https://doi.org/10.1158/1541-7786.mcr-06-0227>.
81. Barry, J.B.; Giguère, V. Epidermal Growth Factor-Induced Signaling in Breast Cancer Cells Results in Selective Target Gene Activation by Orphan Nuclear Receptor Estrogen-Related Receptor α . *Cancer Res.* **2005**, *65*, 6120–6129. <https://doi.org/10.1158/0008-5472.can-05-0922>.
82. Jarzabek, K.; Koda, M.; Kozłowski, L.; Sulkowski, S.; Kottler, M.-L.; Wolczynski, S. The Significance of the Expression of ERR α as a Potential Biomarker in Breast Cancer. *J. Steroid Biochem. Mol. Biol.* **2009**, *113*, 127–133. <https://doi.org/10.1016/j.jsmb.2008.12.005>.
83. Cavallini, A.; Notarnicola, M.; Giannini, R.; Montemurro, S.; Lorusso, D.; Visconti, A.; Minervini, F.; Caruso, M.G. Oestrogen Receptor-Related Receptor Alpha (ERR α) and Oestrogen Receptors (ER α and ER β) Exhibit Different Gene Expression in Human Colorectal Tumour Progression. *Eur. J. Cancer* **2005**, *41*, 1487–1494. <https://doi.org/10.1016/j.ejca.2005.04.008>.
84. Fujimura, T.; Takahashi, S.; Urano, T.; Kumagai, J.; Ogushi, T.; Horie-Inoue, K.; Ouchi, Y.; Kitamura, T.; Muramatsu, M.; Inoue, S. Increased Expression of Estrogen-Related Receptor Alpha (ERR α) Is a Negative Prognostic Predictor in Human Prostate Cancer. *Int. J. Cancer* **2007**, *120*, 2325–2330. <https://doi.org/10.1002/ijc.22363>.

85. Hamidian, A.; von Stedingk, K.; Thorén, M.M.; Mohlin, S.; Pählman, S. Differential Regulation of HIF-1 α and HIF-2 α in Neuroblastoma: Estrogen-Related Receptor Alpha (ERR α) Regulates HIF2A Transcription and Correlates to Poor Outcome. *Biochem. Biophys. Res. Commun.* **2015**, *461*, 560–567. <https://doi.org/10.1016/j.bbrc.2015.04.083>.
86. Zou, C.; Yu, S.; Xu, Z.; Wu, D.; Ng, C.; Yao, X.; Yew, D.T.; Vanacker, J.; Chan, F.L. ERR α Augments HIF-1 Signalling by Directly Interacting with HIF-1 α in Normoxic and Hypoxic Prostate Cancer Cells. *J. Pathol.* **2014**, *233*, 61–73. <https://doi.org/10.1002/path.4329>.
87. Ao, A.; Wang, H.; Kamarajugadda, S.; Lu, J. Involvement of Estrogen-Related Receptors in Transcriptional Response to Hypoxia and Growth of Solid Tumors. *Proc. Natl. Acad. Sci. USA* **2008**, *105*, 7821–7826. <https://doi.org/10.1073/pnas.0711677105>.
88. Sahu, A.; Wang, X.; Munson, P.; Klomp, J.P.G.; Wang, X.; Gu, S.S.; Han, Y.; Qian, G.; Nicol, P.; Zeng, Z.; et al. Discovery of Targets for Immune–Metabolic Antitumor Drugs Identifies Estrogen-Related Receptor Alpha. *Cancer Discov.* **2023**, *13*, 672–701. <https://doi.org/10.1158/2159-8290.cd-22-0244>.
89. Zhao, L.; Li, G.; Meng, F.; Sun, Z.; Liu, J. Cortical and Medullary Oxygenation Evaluation of Kidneys with Renal Artery Stenosis by BOLD-MRI. *PLoS ONE* **2022**, *17*, e0264630. <https://doi.org/10.1371/journal.pone.0264630>.
90. Keppner, A.; Maric, D.; Orlando, I.M.C.; Falquet, L.; Hummler, E.; Hoogewijs, D. Analysis of the Hypoxic Response in a Mouse Cortical Collecting Duct-Derived Cell Line Suggests That Esrra Is Partially Involved in Hif1 α -Mediated Hypoxia-Inducible Gene Expression in mCCDcl1 Cells. *Int. J. Mol. Sci.* **2022**, *23*, 7262. <https://doi.org/10.3390/ijms23137262>.
91. Romero, N.M.; Dekanty, A.; Wappner, P. Cellular and Developmental Adaptations to Hypoxia: A Drosophila Perspective. *Methods Enzymol.* **2007**, *435*, 123–144. [https://doi.org/10.1016/s0076-6879\(07\)35007-6](https://doi.org/10.1016/s0076-6879(07)35007-6).
92. Li, Y.; Padmanabha, D.; Gentile, L.B.; Dumur, C.I.; Beckstead, R.B.; Baker, K.D. HIF- and Non-HIF-Regulated Hypoxic Responses Require the Estrogen-Related Receptor in Drosophila Melanogaster. *PLoS Genet.* **2013**, *9*, e1003230. <https://doi.org/10.1371/journal.pgen.1003230>.
93. Figueiredo, H.; Figueroa, A.L.C.; Garcia, A.; Fernandez-Ruiz, R.; Broca, C.; Wojtusciszyn, A.; Malpique, R.; Gasa, R.; Gomis, R. Targeting Pancreatic Islet PTP1B Improves Islet Graft Revascularization and Transplant Outcomes. *Sci. Transl. Med* **2019**, *11*, eaar6294. <https://doi.org/10.1126/scitranslmed.aar6294>.
94. Shires, S.E.; Quiles, J.M.; Najor, R.H.; Leon, L.J.; Cortez, M.Q.; Lampert, M.A.; Mark, A.; Gustafsson, Å.B. Nuclear Parkin Activates the ERR α Transcriptional Program and Drives Widespread Changes in Gene Expression Following Hypoxia. *Sci. Rep.* **2020**, *10*, 8499. <https://doi.org/10.1038/s41598-020-65438-7>.

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.